

Supporting information for

**Comprehensive analysis of posttranslational protein modifications in aging of
subcellular compartments**

Tim Baldensperger,¹ Michael Eggen,¹ Jonas Kappen,¹ Patrick R. Winterhalter,² Thorsten
Pfaffmann³ and Marcus A. Glomb^{1, *}

¹ Institute of Chemistry, Food Chemistry, Martin-Luther-University Halle-Wittenberg, Kurt-Mothes-Str. 2, 06120 Halle/Saale, Germany

² Clinic for Heart Surgery, Martin-Luther-University Halle-Wittenberg, Ernst-Grube Str. 40, 06120 Halle/Saale, Germany

³ Institute of Physiological Chemistry, Martin-Luther-University Halle-Wittenberg, Hollystr. 1, 06114 Halle/Saale, Germany

* To whom correspondence should be addressed (e-mail marcus.glomb@chemie.uni-halle.de, Tel. +49 345 552-5784, Fax. +49 345 552-7341)

Figure S1. Full-length blotting images of fractionation control.	S-2
Figure S2. Structural formulas of analytes.	S-3
Figure S3. Chromatographic separation of analytes by HPLC-MS/MS.	S-4




Figure S1. Full-length blotting images of fractionation control. Antibodies against cytosolic β -actin (**a**), mitochondrial COX IV (**b**), and histone H3 (**c**) were used.




Figure S2. Structural formulas of analytes.




Figure S3. Chromatographic separation of analytes by HPLC-MS/MS in enzymatic (a) and acid (b) hydrolysates (1 citrulline; 2 *N*^δ-glycoloyl/glyoxylyl lysine; 3 *N*^δ-formyl lysine; 4 *N*^δ-malonyl lysine; 5 *N*^δ-lactoyl/pyruvoyl lysine; 6 *N*^δ-acetyl lysine; 7 *N*^δ-succinyl lysine; 8 *N*^δ-propionyl lysine; 9 methylglyoxal hydroimidazolone; 10 *N*^δ-butyryl lysine; 11 methionine sulfone; 12 methionine sulfoxide; 13 *N*^δ-carboxymethyl lysine; 14 *N*^δ-carboxyethyl lysine; 15 glyoxal hydroimidazolone; 16 furosine; 17 *o*-tyrosine; 18 *o,o*-dityrosine).